


# Container label



The image shows a container label for Folicur 3.6 F. At the top center is the Bayer logo. Below it, the product name "Folicur<sup>®</sup> 3.6 F" is written in large green and black letters. A green banner on the left side of the label indicates "Net Contents: 1 Gallon". The label is divided into two main sections. The left section contains the following text: "For control of specified diseases on peanuts and grasses grown for seed.", "ACTIVE INGREDIENT: Tebuconazole, alpha-(2-(4-chlorophenyl)ethyl)-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol . . . . . 38.7%", "INERT INGREDIENTS: . . . . . 61.3%", "TOTAL: 100.0%", "Contains 3.6 pounds tebuconazole per gallon.", and "EPA Reg. No. 264-752". The right section contains the following text: "STOP - Read the label before use", "KEEP OUT OF REACH OF CHILDREN", "CAUTION", "FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Inside Booklet.", "For PRODUCT USE information Call 1-866-99BAYER (1-866-992-2937)", "For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577", and a vertical code "E004880 UC1 2/22 10/06". At the bottom center, the Bayer CropScience LP address and contact information are listed: "Bayer CropScience LP, P.O. Box 12014, 2 T.W. Alexander Drive, Research Triangle Park, North Carolina 27709, FOLICUR is a registered trademark of Bayer, ©2006 Bayer CropScience".

**BAYER**

# Folicur<sup>®</sup> 3.6 F

**Net Contents:**  
**1 Gallon**

*For control of specified diseases on peanuts and grasses grown for seed.*

**ACTIVE INGREDIENT:**  
Tebuconazole, alpha-(2-(4-chlorophenyl)ethyl)-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol . . . . . 38.7%

**INERT INGREDIENTS:** . . . . . 61.3%

**TOTAL:** 100.0%

Contains 3.6 pounds tebuconazole per gallon.

EPA Reg. No. 264-752

**STOP - Read the label before use**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Inside Booklet.

For PRODUCT USE information Call 1-866-99BAYER (1-866-992-2937)  
For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

E004880 UC1 2/22 10/06

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
FOLICUR is a registered trademark of Bayer.  
©2006 Bayer CropScience

# Container label

NET CONTENTS: 1 GALLON



(01) 0 0785740 13196 1

**FOLICUR® 3.6F Foliar Fungicide**

For control of specified diseases on peanuts and grasses grown for seed.

**ACTIVE INGREDIENT:** Tebuconazole,

alpha-[2-[4-chlorophenyl]ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol ..... 88.7%

**INERT INGREDIENTS:** ..... 61.3%

Contains 3.6 pounds tebuconazole per gallon.

**TOTAL: 100.0%**

EPA Reg. No. 264-752

**Stop - Read the label before use  
KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See Attached Booklet

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-990-2937)

For **MEDICAL AND TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

**FIRST AID**

- IF SWALLOWED:**
  - Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by a poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:**
  - Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15 to 20 minutes.
  - Call a poison control center or doctor for treatment advice.
- IF IN EYES:**
  - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- IF INHALED:**
  - Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
  - Call a poison control center or doctor for further treatment advice.

**Symptoms of Poisoning:** The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**  
Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

**FOR ADDITIONAL PRECAUTIONARY STATEMENTS:** See attached label booklet: Personal Protective Equipment (PPE),

User Safety Recommendations, and Environmental Hazards

**DIRECTIONS FOR USE:** See attached booklet.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

**Note To Physician:** No specific antidote. Treat symptomatically.

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
FOLICUR is a registered trademark of Bayer.  
©2006 Bayer CropScience

6004880 021210A 08/06

# Container label

## Warranty statement

### **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

# Web distributed labeling

## Peanut Use Directions – geographic restrictions

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Peanut	<b>SOILBORNE:</b> Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) <b>FOLIAR:</b> Early leaf spot Late leaf spot Leaf rust Web blotch (Phoma) Pepper spot (Leptosphaerulina)	7.2 fl oz per acre
	<p><b>FOUR-APPLICATION SPRAY PROGRAM:</b> Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of FOLICUR 3.6 F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F.</p> <p><b>LEAF SPOT ADVISORY SCHEDULE:</b> For control of soilborne diseases in an advisory schedule, apply FOLICUR 3.6 F in the first advisory spray in July and continue FOLICUR 3.6 F applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.</p>	
<p><b>GENERAL DIRECTIONS:</b> For optimum control of the specified soilborne diseases, four consecutive applications of FOLICUR 3.6 F must be made at 14-day intervals.</p> <p>A maximum of 28.8 fluid ounces of FOLICUR 3.6 F may be applied per crop season. FOLICUR may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.</p> <p>FOLICUR 3.6 F is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with FOLICUR 3.6 F as a leaf spot resistance management strategy. A spray surfactant is not necessary when FOLICUR 3.6 F is tank mixed with chlorothalonil. Mixing or alternating FOLICUR 3.6 F with other DMI fungicides may lead to resistance.</p> <p>FOLICUR 3.6 F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of FOLICUR 3.6 F against the root and pod rots.</p> <p>Use FOLICUR 3.6 F in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
<b>Timing of FOLICUR 3.6 F Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot</b>		
<b>Spray Program</b>	<b>FOLICUR 3.6 F Application No.</b>	<b>Chlorothalonil Application No.</b>
7 applications	3, 4, 5 and 6	1, 2 and 7

# Web distributed labeling

## **OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

## **ROTATIONAL CROPS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.